Fact Sheet - Lao

ຫ່ວງ IUD ແບບປ່ອຍຮໍໂມນ (MIRENA®)

The Hormone Releasing IUD (Mirena®)



ຫ່ວງ IUD ແບບປ່ອຍຮໍໂມນ ແມນຫຍັງ?

ຫ່ວງ IUD ແບບປ່ອຍຮໍໂມນ ຊະນິດ ດຽວທີ່ມີຈຳໜ່າຍຢູ່ໃນອອສເຕຣເລັຍ ເອີ້ນວ່າ Mirena®. ມັນແມ່ນເຄື່ອງ ປລາສຕິກນ້ອຍໆອັນໜຶ່ງທີ່ມີຮູບຮ່າງຄ້າຍ ໂຕ 'T' ທີ່ມີຮໍໂມນຊະນິດໜຶ່ງເອີ້ນວາ



ລີເວີໂນແຈສໂຕຣ ຢູ່ໃນກ້ານຂອງມັນ. ຫ່ວງຈະເອົາໃສ່ໄວ້ມີດລູກ ເພື່ອປ້ອງກັນບໍ່ໃຫ້ຕັ້ງທ້ອງ. ມັນມີສາຍໄນລອນບາງໆຕິດຢູ່ແລະເມື່ອ IUD ຖືກໃສ່ເຂົ້າທີ່ແລ້ວ, ສາຍນັ້ນກໍຈະຍ່ອນອອກມາທາງປາກມືດ ລູກ (ບອນສຸດຂອງມຶດລູກ) ຜານຊອງຄອດ. ຫວງ IUD ແບບປອຍຮໍ ໂມນຈະຢູ່ໃນທີ່ຂອງມັນໄດ້ຮອດຫ້າປີ.

ຫວງ IUD ແບບປອຍຮໍໂມນ ເຮັດວຸງກແບບໃດ?

ຮໍໂມນຈະເຮັດໃຫ້ນໍາເມືອກໃນປາກມິດລູກໜາຂຶ້ນ ສະນັ້ນ ຕົວ ເຊື້ອສະເປີມຈຶ່ງບໍ່ສາມາດເຂົ້າໄປທາງໃນມືດລູກໄດ້ແລະ ນອກນັ້ນ ມັນຈະເຮັດໃຫ້ຜະໜັງມິດລູກບາງລົງ. ບາງເທື່ອມັນກໍອາດເຮັດໃຫ້ ຮວຍໄຂຢຸດປອຍໄຂອອກມາ ເຊັ່ນກັນ.

ມັນໃຊ້ໄດ້ຜິນດີປານໃດ?

ຫ່ວງ IUD ແບບປ່ອຍຮໍໂມນ ແມ່ນວິທີໜຶ່ງທີ່ໄດ້ຜົນດີຫຼາຍໃນ ການປ້ອງກັນບໍ່ໃຫ້ຕັ້ງທ້ອງ (ໄດ້ຜົນ 99.8%). ນີ້ໝາຍຄວາມວ່າ ມີແມ່ຍິງນ້ອຍກວ່າ ໜຶ່ງຄົນໃນຈຳນວນແມ່ຍິງແຕ່ລະຮ້ອຍຄົນທີ່ໃຊ້ ວິທີຄຸມກຳເນີດແບບນີ້ ທີ່ຈະຕັ້ງທ້ອງ ໃນປີໜຶ່ງ.

ແມນໃຜໃຊ ຫວງ IUD ແບບປອຍຮໍໂມນ ໄດ?

ແມ່ຍິງສ່ວນໃຫຍ່ສາມາດໃຊ້ IUD ໄດ້, ຮວມທັງແມ່ຍິງ ທີ່ບໍ່ເຄີຍມີລູກມາກ່ອນ, ແລະແມ່ຍິງທີ່ໃຫ້ລູກກິນນົມແມ, ແຕ່ວ່າໃຫ້ຖາມທ່ານໝໍເບິ່ງ ວ່າມັນເໝາະສົມກັບທ່ານຫຼືບໍ່. ມັນຈະເປັນການດີເຊັ່ນກັນ ຕໍ່ແມ່ຍິງທີ່ປະຈຳເດືອນມາແຮງ ເພາະ ວ່າມັນມີແນວ ໂນ້ມວ່າຈະເຮັດ ໃຫ້ປະຈຳເດືອນສັ້ນລົງແລະເບົາ ຂຶ້ນ (ແລະສ່ວນໃຫຍ່ອາດເຮັດໃຫ້ປະຈຳເດືອນຢຸດໄປເລີຍ). ອັນນີ້ ຈະບໍ່ເປັນອັນຕະລາຍ ຫຍັງຕໍ່ຮ່າງກາຍຂອງທ່ານ.

ແມ່ນໃຜບໍ່ຄວນໃຊ[້] ຫ່ວງ IUD ຊະນິດນີ້?

ທ່ານບໍ່ຄວນໃຊ້ຫ່ວງນີ້ຖ້າທ່ານອາດຈະຕັ້ງທ້ອງແລ້ວ. ມີອາການທາງ ສຸຂະພາບບາງຢ່າງເຊິ່ງທ່ານມີທີ່ອາດເຮັດໃຫ້ທ່ານໃຊ້ຫ່ວງ IUD ີ່ ໂປຣແຈສໂຕເຈີນບໍ່ໄດ້. ມັນເປັນສິ່ງສຳຄັນຫຼາຍທີ່ທ່ານຕ້ອງປຶກສາ

ກັບທ່ານໝໍຂອງທ່ານເພື່ອໃຫ້ໝັ້ນໃຈວ່າເປັນການປອດໄພສຳລັບທ່ານ ທີ່ຈະໃຂ້ຫ່ວານີ້.

ມີຜືນຂາງຄຽງຫຍັງບໍ?

ແມ່ຍິງອາດເປັ້ນເລືອດບໍ່ສະມ່ຳສະເໝີເປັນກະປິດກະປອຍໃນ ສາມສີ່ເດືອນທໍາອິດຫຼັງຈາກໄດ້ສອດ IUD ແບບປ່ອຍຮໍໂມນນີ້. ຕາມປົກກະຕິ ການເປັນເລືອດແບບນີ້ຈະເຊົາແລະປະຈຳເດືອນກໍຈະ ສັ້ນລົງແລະເບົາຂຶ້ນແລະສ່ວນຫຼາຍ ປະຈຳເດືອນກໍຈະບໍ່ມາເລີຍ. ແມ່ຍິງບາງຄືນອາດມີຜົນຂ້າງຄຸງທາງຮໍໂມນ, ເຊັ່ນ ທ້ອງອືດຫຼືຜິວພັນປຸ່ງນແປງ, ແຕ່ວ່າອັນນີ້ບໍ່ຄ່ອຍເກີດຂຶ້ນເພາະວ່າປະ ລິມານຂອງຮໍໂມນນັ້ນມີນ້ອຍຫຼາຍ.

ມີເປີເຊັນເລັກນ້ອຍວ່າຈະເຮັດໃຫ້ເກີດການຕິດເຊື້ອແຖບກະໂທ້ຍ ໃນເວລາທີ່ສອດ IUD.

ທ່ານໝໍແລະພະຍາບານຈະອະທິບາຍໃຫ້ທ່ານຟັງກ່ຽວກັບຄວາມສ່ຽງ ຢ່າງອື່ນໃນການສອດ IUD.

ທານຈະເອົາຫ່ວງ IUD ແບບປອຍຮໍໂມນໄດ້ຢ່າງໃດ?

ທ່ານຕ້ອງໄປຫາທ່ານໝໍຫຼືໄປທີ່ຄລີນິກວາງແຜນຄອບຄົວ. ທ່ານອາດຖືກຂໍໃຫ້ໄປສອງເທື່ອ. ເທື່ອທຳອິດ ທ່ານໝໍຈະຖາມກ່ຽວ ກັບສຸຂພາບຂອງທານແລະອະທິບາຍໃຫ້ທານຟັງກຸງວກັບຫວງ IUD. ທ່ານຈະໄດ້ຮັບການກວດຊ່ອງຄອດ/ກະໂທ້ຍ, ແລະອາດກວດເບິ່ງວ່າ ມີການຕິດເຊື້ອໃນຊ[່]ອງຄອດ ຫຼືບໍ່. ຄັ້ງທີສອງເພິ່ນຈະສອດ IUD ໃຫ[້]. ທ່ານໝໍຫຼືພະຍາບານຈະອະທິບາຍຂັ້ນຕອນການສອດນີ້ແກ່ທ່ານ. ຈະໃຊ້ເວລາປະມານ 10 ນາທີແລະແມ່ຍິງບາງຄົນອາດຮູ້ສຶກບໍ່ສະ ບາຍໃນຂະນະທີ່ບາງຄົນອາດຮູ້ສຶກເຈັບ. ເປັນການດີທີ່ຈະກິນຢາແກ້ ປວດແບບທຳມະດາກອນເຮັດການສອດ. ທານບໍ່ຈຳເປັນຕ້ອງອີດອາ ຫານກ່ອນເຮັດການສອດ IUD, ຄວາມຈິງແລ້ວແມ່ນເປັນການດີທີ່ ຈະກິນກ່ອນໜ້ານັ້ນ. ທ່ານອາດຮູ້ສຶກເປັນວິນໃນລະຫວ່າງທີ່ສອດໃສ່ຫຼື ້ຫຼັງຈາກນັ້ນແລະທ່ານຈະຕ້ອງພັກຜ່ອນລະຍະໜຶ່ງກ່ອນອອກຈາກຄລີ ນິກ. ທ່ານຄວນກະເວລາໄວ້ປະມານຊື່ວໂມງໜຶ່ງຢູ່ຄລີນິກ.

ຫວງ IUD ແບບປອຍຮໍໂມນຖືກເອົາອອກແບບໃດ?

ທ່ານຕ້ອງໄປຫາໝໍຫຼືຄລີນິກວາງແຜນຄອບຄົວເພື່ອເອົາອອກ. ທ່ານ ໝໍຈະໃຊ້ເຄື່ອງມືພິເສດເພື່ອເອົາຫ່ວງ IUD ອອກ ໂດຍດຶງສາຍເຊືອກ ຄ່ອຍໆ. ອັນນີ້ຈະໃຊ້ເວລາແຕ່ພູເງສອງສາມນາທີ. ແມ່ຍິງບາງຄົນອາດ ຮູ້ສຶກບໍ່ສະບາຍແຕ່ບາງຄົນກໍບໍ່ຮູ້ສຶກ ຫຍັງເລີຍ.

www.fpnsw.org.au | talkline 1300 658 886 | bookshop

ບໍລິການປິ່ນປົວ&ໃຫ້ ຂໍ້ມູນ | ສຶກສາ ຝຶກງານ | ຄົ້ນຄວ້າ | ພັດທະນາສາກິນ

ພະແນກວາງແຜນຄອບຄົວ ຣັດ NSW ແມ່ນອີງການທີ່ບໍ່ຫວັງຜົນກຳໄລ ໄດ້ຮັບການອຸປະຖຳໂດຍກະຊວງສາທາຣະນະສຸກ ຣັດ NSW.





ສິ່ງທີ່ຄວນຈຳ

ໃຫ້ຄຳເບິ່ງທາງໃນຊ່ອງຄອດດ້ວຍນິ້ວມືເພື່ອກວດເບິ່ງເຊືອກ ທຸກໆເດືອນ ຫຼັງປະຈຳເດືອນຢຸດ ເພື່ອໃຫ້ໝັ້ນໃຈວ່າ IUD ຍັງຢູ່ຄືເກົ່າ. ຖ້ຳທ້ານມີອາການຜິດປົກກະຕິໃດໆ ຮວມທັງມີນ້ຳລົງຂາວອອກຈາກຊ່ອງຄອດ, ເຈັບທ້ອງນ້ອຍຫຼືເຈັບໃນ ລະຫວ່າງຮ່ວມເພດ, ໃຫ້ໄປຫາໝໍຂອງທ່ານທັນທີ. ຖ້າທ່ານກັງວົນວ່າທ່ານອາດຕັ້ງທ້ອງແລ້ວ, ໃຫ້ໄປຫາທ່ານໝໍຫຼື ຄລີນິກວາງແຜນຄອບຄົວເພື່ອກວດເບິ່ງວ່າໄດ້ຕັ້ງທ້ອງຫຼືບໍ່. ແຕ່ວ່າ ຂໍໃຫ້ຈຳໄວ້ວ່າ, ປະຈຳເດືອນທ່ານອາດບໍ່ມາເມື່ອທ່ານໃຊ້ IUD ແບບປ່ອຍຮໍໂມນ. ຖ້າທ່ານ, ຫຼືຄູ່ຄອງຂອງທ່ານ, ເຄີຍມີເຊັກສ໌ແບບຊື່ວຄັ້ງຊື່ວຄາວ, ຫຼືວ່າທ່ານໄດ້ຄູ່ນອນຄົນໃໝ່, ຕ້ອງໃຊ້ຖຶງອະນາໄມທຸກຄັ້ງ ຈົນກວ່າ ທັງສອງຄົນໄດ້ຮັບການກວດຫາເຊື້ອກາມະໂຣກ (STIs)ແລ້ວ. ໃຫ້ຂູງນບັນທຶກວັນທີ່ຕ້ອງປຸ່ງນຫ່ວງເພາະວ່າຫ່ວງ IUD ບໍ່ຄວນຖືກປະ ໄວ້ເກີນກວ່າເວລາທີ່ແນະນໍ້າ.

ຖ້າຕ້ອງການຂໍ້ມູນເພີ້ມເຕີມ

- ໃຫ້ຕິດຕໍ່ສາຍສິນທະນາຂອງພະແນກວາງແຜນຄອບຄົວແຫ່ງລັດ NSW ທີ່ເລກ ໂທ 1300 658 886 ຫຼືເຂົ້າໄປເບິ່ງ www.fpnsw.org.au/talkline
- NRS (ສຳລັບຄົນພິການຫູ) 133 677ຫຼືໄປທີ່ຄລີນິກວາງແຜນຄອບຄົວໃກ້ບ້ານທ່ານ

www.fpnsw.org.au | talkline 1300 658 886 | bookshop

ບໍລິການປິ່ນປົວ&ໃຫ້ ຂໍ້ມູນ | ສຶກສາ ຝຶກງານ | ຄົ້ນຄວ້າ | ພັດທະນາສາກິນ ພະແນກວາງແຜນຄອບຄົວ ຣັດ NSW ແມ່ນອີງການທີ່ບໍ່ຫວັງຜົນກຳໄລ ໄດ້ຮັບການອຸປະຖຳໂດຍກະຊວງສາທາຣະນະສຸກ ຣັດ NSW.






THE HORMONE-RELEASING IUD (MIRENA®)



What Is The Hormone-releasing IUD?

The only Hormone-releasing IUD available in Australia is called Mirena®. It is a small plastic device in the shape of a 'T' with a hormone called levonorgestrel in its stem. It is placed inside the uterus to prevent pregnancy. It has a fine nylon string attached and when the IUD is in place, the string comes out through the cervix (the end of the uterus) into the vagina. The Hormone-releasing IUD can stay in place for up to five years.



How Does Hormone-releasing IUD Work?

The hormone makes the mucus in the cervix thicker so that sperm cannot get into the uterus and it also thins the lining of the uterus.. It can also sometimes stop your ovaries from releasing an egg.

How Well Does It Work?

The hormone-releasing IUD is a very effective method of preventing a pregnancy (99.8% effective). This means that fewer than one woman in every hundred using this method of contraception for a year would become pregnant.

Who Can Use Hormone-releasing IUD?

Most women are able to use an IUD, including women who haven't had a baby before, and women who are breastfeeding, but ask your doctor if it is suitable for you. It is also useful for women who have heavy periods because it tends to make periods shorter and lighter (and can often stop periods altogether). This is not harmful in any way to your body.

Who Should Not Use This Type Of IUD?

You should not use it if you could be pregnant. There are some medical conditions that you may have which could prevent you from using the Progestogen IUD. It is very important to talk with your doctor to be sure it is safe for you to use it.

Are there any side effects

Women may have irregular bleeding and spotting in the first few months after the hormone-releasing IUD is inserted. Bleeding usually settles down and periods become shorter and lighter and often women will have no periods at all.

Some women may experience hormonal side effects, such as bloating or skin changes, but this is uncommon because the dose of hormone is so small.

There is a small chance of getting a pelvic infection at the time of the IUD insertion.

The doctor or nurse will tell you about other risks of having the IUD inserted.

How Do You Get The Hormone-releasing IUD?

You need to go to a doctor or Family Planning Clinic. You may be asked to make two visits. On the first visit the doctor asks questions about your health and tells you about the IUD. You will have a vaginal/pelvic examination, and possibly a test for vaginal infection. On the second visit the IUD is inserted. Your doctor or nurse will explain the procedure to you. It takes about 10 minutes and some women may find the experience uncomfortable while others may find it quite painful. It is a good idea to take simple pain tablets before the procedure. You do not have to fast before having the IUD put in, in fact it is a good idea to eat beforehand. You may feel faint during or after the insertion and you will probably need to rest for a while before you leave the clinic. You should allow about an hour to be at the clinic.

www.fpnsw.org.au | talkline 1300 658 886 | bookshop

clinical services & information | education & training | research | international development Family Planning NSW is a not-for-profit organisation funded by the NSW Ministry of Health





How Is The Hormone-releasing IUD Taken Out?

You need to go to a doctor or Family Planning Clinic to have it taken out. The doctor uses a special instrument to remove the IUD by gently pulling on the string. This only takes a couple of minutes. Some women find it a little uncomfortable and some don't feel much at all.

Things To Remember

Feel inside your vagina with your finger to check for the string each month after a period to be sure the IUD is in place.

If you have any unusual symptoms including a discharge from your vagina, pain low in your abdomen or deep pain during intercourse, see your doctor straight away.

If you are concerned that you could be pregnant, see your doctor or clinic for a pregnancy test. But remember, you can miss periods with the hormone-releasing IUD.

If you, or your partner, ever have casual sex, or if you have a new sexual partner, use a condom every time until you both have been checked for sexually transmissible infections (STIs).

Keep a record of the date for replacement because an IUD should not stay in for longer than the recommended time.

For Further Information

- Contact the Family Planning NSW Talkline on 1300 658 886 or go to www.fpnsw.org.au/talkline
- NRS (for deaf) 133 677
- Or visit your nearest Family Planning clinic

Reviewed: June 2014 | FPNSW 06/14

The information in this Factsheet has been provided for educational purposes only. Family Planning NSW has taken every care to ensure that the information is accurate and up-to-date at the time of publication. Individuals concerned about any personal reproductive or sexual health issue are encouraged to seek advice and assistance from their health care provider or visit an Family Planning NSW clinic.



 $\label{lem:clinical services & information | education \& training | research | international development \\ \textit{Family Planning NSW} is a not-for-profit organisation funded by the NSW Ministry of Health \\ \\$

